This form is for people who have a moderately to severely weakened immune system who have already received 2 doses of Pfizer or Moderna COVID-19 vaccine.

If you meet the <u>criteria</u> outlined by the CDC, it is recommended that you get a 3rd dose of an mRNA (Pfizer or Moderna) vaccine at least 28 days after your 2nd dose of vaccine. When possible, you should receive the same vaccine. For example, if you got a series of Pfizer vaccine, try to get a Pfizer vaccine for your 3rd dose. Talk to your doctor about the need to get a 3rd dose of COVID-19 vaccine and the best timing of a 3rd dose.

- ☐ I attest that I am immunocompromised and am eligible for a third dose of vaccine based on the criteria below.
 - Receiving active cancer treatment for tumors or cancers of the blood;
 - Received an organ transplant and am taking medicine to suppress my immune system;
 - Received a stem cell transplant within the last 2 years or am taking medicine to suppress the immune system;
 - Have moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome);
 - Have advanced or untreated HIV infection;
 - Receiving active treatment with high-dose corticosteroids or other drugs that suppress my immune response

Signature:	
Name:	Date:

It is important to continue to protect yourself after you get a 3rd dose of vaccine. This includes wearing a well-fitting mask, maintaining physical distance, avoiding crowded places or spaces with poor air flow, and washing hands often. Consider "double masking" (wearing a cloth face mask over surgical mask) or an N95 respirator for a higher level of protection.

For more information, see the <u>Moderna</u> or <u>Pfizer</u> Fact Sheet for Recipients and Caregivers and the CDC webpage <u>COVID-19 Vaccines for Moderately to Severely Immunocompromised People</u>.

If you do not have access to a printer, you can write the attestation by hand.

